

SUBJECT INDEX

A

- A stars, peculiar, 432-35
- Abundances
 - in interstellar medium, 525-31
 - molecular, in stars, 92-96, 102-3
 - solar, 398-99
- Accretion disk in active galaxies, 172, 178, 194, 196-98
- Accretion phase of stellar evolution, 52
- Active galactic nuclei, 171-203.
 - See also Seyfert galaxies, Active galaxies
 - black holes in, 172
 - polarization in, 183-84
- Active galaxies, 171-203
 - cloud models for broad-line region of, 186-96
 - noncloud models for broad-line region of, 196-98
 - photoionization in, 173, 182, 185-90
- Akn 120, 179, 188
- Akn 564, 177
- Alpha Aquarii, 384, 388, 398, 412
- Alpha Aurigae, 404
- Alpha Bootes, 113-14, 390, 400
- Alpha Cygni, 413
- Alpha Eridani, 522-23
- Alpha Herculis, 116, 379, 390
- Alpha Orionis, 114, 117, 388, 390, 401-2
- Alpha Scorpii, 390
- Alpha Tauri, 390
- Alpha Trianguli Australis, 387-88, 394
- Aperture synthesis, 132-33, 138, 147-49, 162-63
- Atomic clock, 538-40, 548-49

B

- B stars, interstellar reddening of, 14-16
- Bayesian method, 137-38, 141, 146-47
- Becklin-Neugebauer object, 81
- Beta Aquarii, 384, 412
- Beta Geminorum, 390
- Beta Ursae Minoris, 390
- Binary systems
 - mass-loss rates from, 368, 378, 403-5

- pulsars in, 286, 292, 303-7, 323, 537, 554-67, 572-73
- RS Canum Venaticorum-type, 400, 404-5, 414
- Wolf-Rayet stars in, 362, 364, 368, 370
- Black holes
 - in active galactic nuclei, 172
 - in quasars, 189-90
 - and supernova explosions, 218, 226, 361

C

- Carbon stars, 93-96, 102, 105-10, 113-14, 116
- Cassiopeia A, 220, 223, 228, 347-48
- Centaurus A, 468-69
- Charge-coupled devices, 21, 255-83
 - astronomical applications of, 280-81
 - charge-transfer efficiency in, 269-71
 - drift scanning in, 278-79
 - effect of cosmic-ray events on, 268
 - flat-fielding procedures for, 276-79
 - intrapixel nonuniformity in, 279
 - luminescence effects on, 268
 - photon detection in silicon in, 265-67
 - principles of, 256-61
 - readout noise in, 273-76
 - readout procedures in, 271-76
 - thermal effects on, 267-68
 - used for detecting interstellar gas, 510
 - used as X-ray detectors, 282
- Circumstellar envelopes, 115-20, 402-3, 407, 413
- Circumstellar lines, 385, 392, 408
- Circumstellar shells, 337, 339-40, 345-47, 382, 388
- Color-magnitude diagrams, 437, 444-53
- Cool stars, 377-420. See also G stars, K stars, M stars
 - chromospheres of, 378-95, 399-402, 410
 - mass loss from, 377-420
 - dwarf, 378, 380-81, 384-86, 391, 399

- giant, 378-79, 382-88, 390-91, 393, 403-4, 407-12
- supergiant, 378-79, 381-88, 390-91, 400-4, 408, 412-13
- mass-loss rates from, 337-38
- molecular opacities of, 96-101
- spectral classification of, 104, 106-11
- Copernicus satellite observations, 329, 500, 511, 516, 518, 520, 524-25, 530
- Coronal mass ejections, 395
- Cosmic microwave background radiation, 530-31
- Cosmic-ray events, effect on CCDs of, 268
- Cosmic rays, 192, 461, 463, 472, 479
- Crab Nebula pulsar, 208, 293-95, 299, 318-19, 539
- CS stars, 93-94

D

- Deuterium, interstellar, 529
- Doppler shifts in solar spectrum, 35-39
- Dynamo theory, 460, 481-86

E

- Einstein telescope observations, 132
- Emission nebulae, 50-52. See also H II regions
- Eta Carinae, 357

F

- F stars, 434-35
- Faraday rotation, 460-67, 469-80, 487
- Fornax system, 422, 426-27, 435
- Fourier transform, 127-33, 159, 165

G

- G stars
 - giant, 379, 382, 404
 - supergiant, 383-85, 388, 412-13
- Galactic bulge, stellar population of, 19-22

- Galactic center, 19
 Galactic halo, 479-80, 487-88
 Galactic structure, star counts and, 577-611
- Galaxies
 colors of, 18
 magnitudes of, 16-17
 molecules as population indicators in, 110-11
 stellar populations of, 21
 Galaxies, barred spiral, 488-89
 Galaxies, dwarf, 489
 Galaxies, edge-on, 468, 478-80
 Galaxies, elliptical, 18, 421, 425-28, 435-37
 Galaxies, irregular, 469, 477
 Galaxies, peculiar, 469
 Galaxies, spiral, 425, 428
 magnetic fields in, 459-97
 Sd, 474-76
 supernovae in, 206-7
- Galaxy
 black hole in center of, 172
 collapse of, 453-55
 distribution of stars in, 578-83, 585, 596-609
 formation of, 421-23, 453-55
 interstellar medium in, 50-52, 499-535
 interstellar reddening in, 14-16
 magnetic field in, 172, 467-68, 477-78, 487-88
 magnetic halo in, 479-80, 487-88
 pulsar population in, 297-303
 supernovae frequency in, 206
 thick disk of, 602-4
- Galaxy, disk of
 cold-cloud interstellar medium in, 513-15
 hot gas component of interstellar medium in, 518-21
 warm-gas interstellar medium in, 516-18
- Galaxy model, 579-96
 galactic bulge in, 591, 607-9
 globular cluster feature in, 594, 600-1
 kinematics in, 607-8
 massive halo in, 592, 605-6
 Population I disk stars in, 589-93
 luminosity functions of, 592-93, 601, 604
 scale lengths and scale heights of, 595-96, 602
 Population II spheroidal stars in, 589-91, 593-94
 luminosity functions of, 593-94
- scale lengths and scale heights of, 595-96, 602
- Gamma Cassiopeiae, 530
 General relativity, 537-75
 Globular cluster feature of Galaxy model, 594, 600-1
 Globular clusters, 421-28, 436, 515
 ages of, 446, 449-52
 asymptotic giant branch of, 405-7, 453
 colors of, 14-15
 distances of, 447-48
 luminosity functions of, 453, 593-94
 luminosity normalization of, 448-49
 magnetic fields in, 469
 main sequences of, 429-35, 446-53
 mass loss in stars in, 405-10
 stellar evolution and HR diagram of, 429-32
- Gravitation, alternative theories of, 566-67
 Gravitational rotation, 558-60
 Gravitational-wave background, 569-73
 Gravitational waves, pulsar timing and, 567-73
 Groombridge 1830, 438
 G320.4 - 1.2, 294
- H
- H II regions
 blister-type, 77-79, 82-83
 compact, 68, 72-76
 E-type, 50-52, 76
 formation phase of, 69-70
 ionization and recombination in, 59-61, 71-72
 magnetohydrodynamics in, 55-57
 radiation transfer in, 57-59
 S-type, 51-52, 69-83
 temperatures in, 54-63
 ultracompact, 73, 76, 81-82
- H II regions, evolution of, 49-87
 boundary conditions in, 63-68
 champagne phase of, 76-80, 82-83
 expansion phase of, 70-71, 76-80
 microphysical processes in, 55, 64-68
 recombination phase of, 71-72, 80-81
- HD 19557, 95
 HD 19945, 433
 HD 72127A,B, 524
- HD 84903, 410
 HD 93521, 518
 HD 140283, 433
 HD 165195, 408
- HR diagram, 423, 435
 asymptotic giant branch of, 405-7, 453
 main-sequence position in, 430-32, 437-46
 mass-loss effects on, 329, 341-44, 355, 369, 390-95
- HR diagram of globular clusters, 429-32
 HR diagram of luminous stars, 348-53
 HR 1040, 413
 HR 1099, 400, 404
 HR 8752, 413
 Hubble constant, 243
 Hubble-Sandage objects, see Stars, luminous blue variable
- Hyades cluster stars, 386-87
- I
- IC 342, 475-76
- Image restoration, 127-70
 in aperture synthesis mapping, 132-34, 138, 147-49, 162-63
 need for nonlinearity in, 133, 151, 163-66
 in single aperture case, 129-32, 149-50
 superresolution in, 133-34, 144, 154, 165
- Image restoration methods
 a priori information in, 128, 133-34, 140-41, 147, 167
 Clean algorithm for, 128, 152, 162-64, 166-67
 Maximum Entropy Method, 127-70
 positivity-enforcing algorithms for, 165-66
- Infrared observations
 of B stars, 15
 of Galactic bulge, 19-21
 Infrared sources, 52, 75
 Infrared spectra of cool stars, 93-96, 104-10, 114, 117
 Interferometric devices for interstellar gas studies, 508-10
 Interferometric positions of pulsars, 552-53
 Interferometry, image restoration in, 127-29, 132-34, 159-60

- noise-free, 142-46
- with noisy data, 146-49
- Intergalactic magnetic field, 492-93
- Interstellar deuterium, 529-30
- Interstellar gas
 - CCDs used in measuring, 510
 - interferometric devices for measuring, 508-10
 - in local region, 502-3, 521-23
 - optical instruments for measuring, 508-11
 - techniques for measuring, 503-8
 - ultraviolet instruments for measuring, 511-13
- Interstellar lithium, 529-30
- Interstellar magnetic fields, 459-67, 487-92
- Interstellar medium, 49-52, 499-535
 - abundances in, 525-31
 - cold clouds of the, 501-2, 513-15, 525-28
 - enrichment by mass loss, 358-61
 - hot coronal gas in, 501, 503, 518-23
 - and pulsar arrival times, 539-40
 - and supernova remnants, 503, 523-25
 - warm gas of the, 501-3, 516-18, 525-28
- Interstellar reddening, 14-16
- IUE observations, 94, 180, 191, 329, 334, 390, 500, 511, 516, 520
- K
- K stars
 - dwarf, 606
 - giant, 20, 108-9, 113-14, 383
 - supergiant, 383-84
- Kleinmann-Low region, 75, 82
- L
- Lambda Andromedae, 382, 404
- Large Magellanic Cloud, 288, 353, 365, 406, 469, 477
- Liners, 65, 68, 176-77
- Lithium, interstellar, 529-30
- M
- M stars
 - dwarf, 93-94, 105, 109-11, 118
 - emergent fluxes in, 103-5
 - giant
 - in bulge population, 20-22
 - mass loss in, 381-82
 - molecular spectra of, 94-96, 102-5, 108-10, 112, 114, 116-17
 - metal-deficient, 110-11, 381
 - spectral classification of, 93-96, 106-11
 - subdwarf, 110-11
 - supergiant, 109, 114, 117, 119
- Magellanic Clouds, 511. See also Large Magellanic Cloud and Small Magellanic Cloud
- Magnetic field
 - bisymmetric-spiral, 460, 465-67, 470-78, 480, 484-94
 - in Galaxy, 172, 460-64, 467-68, 477-78, 487-88
 - dynamo theory of, 481-86
 - intergalactic, 492-94
 - of neutron stars, 542
 - ring-configuration, 460, 465, 467, 470-71, 480-84, 486, 489-90, 492-93
- Magnetic fields
 - galactic, 492-93
 - global, 487-88
 - interstellar, 459-67, 487-92
 - in pulsars, 294-95, 297, 306, 321-22
 - solar, 24, 26, 28, 31, 44-46, 395
 - in spiral galaxies, 459-97
- Magnetic halo, 478-80, 487-88
- Magnetohydrodynamics, 55-57, 459-60, 478, 490
- Main-sequence position, 437-46, 448-53
- Mass loss
 - from cool stars, 377-420
 - observational constraints on theory of, 413-15
 - radio-frequency determinations of, 388-90
 - detected from spectroscopic observations, 378-90
 - from dwarf stars, 395-400
 - effects on HR diagram of, 329, 341-44, 348-53, 355, 369, 390-95
 - from globular cluster stars, 405-10
 - from massive stars, 329-75
 - coronal theory of, 334, 337
 - fluctuation theory of, 334, 338
 - hot-cool star connection in, 338
 - radiation pressure theory of, 333-34, 337
 - from metal-deficient stars, 352-53, 405-10
 - in Population I stars, 380-81, 386, 390-95, 406
 - from the Sun, 378-79, 384, 386, 395-99
 - from supergiants, 337-38, 344-47, 351-53, 355-56, 366-67
- Mass-loss rates
 - for early-type stars, 333-37, 352
 - for late-type stars, 337-38, 353, 378-90, 400-3
 - for OB supergiants, 333-37, 350-51
 - parameterization of, 410-13
 - radio techniques for determining, 334, 388-90
 - UV techniques for determining, 334, 385-88
 - velocities of, 378-90
 - X rays used in determining, 393-95
- Massive stars
 - galactic distribution of, 366-69
 - as supernova precursors, 206-12, 226-28, 347-48
- Massive stars, evolution of, 329-75
 - at constant mass, 330-33, 347, 355
 - convective overshooting in, 353-55, 368, 406
 - core He-burning phase, 331, 341-47, 355
 - core and shell H-burning phases of, 331, 339-47, 354
 - effects of mass loss on, 339-48
 - effects of metallicity on, 352-53, 366-67
- Maximum entropy images, properties of, 141-54
- Maximum Entropy Method of image restoration, 127-70
 - a priori information in, 140-41
 - algorithms for, 154-60
 - default approach to controlling, 151-52, 161-62, 166
 - least-squares, 147-53, 156, 167
 - nonlinearity in, 143-44, 149
 - for polarization mapping, 159-61
 - resolution variation in, 152

- in single-aperture case, 149–50
- for spectral imaging, 161–62
- Mira variables, 95, 114–17, 402
- Molecular clouds, 50–54, 77–80, 82, 500–1
- Molecular opacities, 96–105, 118
- Molecules
 - fluorescent lines of, 115
 - thermodynamical properties of, 91–92
- Molecules in circumstellar envelopes, 115–20
- Molecules in stars, 89–125
 - abundances of, 92–96, 102–3, 113
 - as population indicators, 110–11
- Mrk 279, 177
- Mrk 359, 177
- Mrk 876, 177
- Mrk 926, 177
- Mrk 1014, 185
- MS stars, 93, 106
- Mu Cassiopeiae, 441
- M 3, 429, 435, 442–43, 449–51, 600
- M 5, 435
- M 13, 430, 435, 453, 594, 600
- M 15, 442, 451–52
- M 22, 515
- M 31, 19, 427, 429, 435–36, 469–70, 481, 489
- M 32, 18, 426–27, 435
- M 33, 471–72
- M 51, 472–73
- M 67, 436
- M 71, 407
- M 81, 469, 473–74, 487
- M 82, 469
- M 83, 476, 488–89
- M 8/, 166, 425
- M 92, 429–30, 435, 442–43, 449, 452, 594, 600
- N
- N stars, 94, 107, 109
- Nebulae
 - bipolar, 50
 - emission, see Emission nebulae
- Neutrino transport model, 213–14, 217–19
- Neutron stars, 205, 208, 218, 285, 293–95, 303–7, 316–17, 323, 542, 560, 573
- NGC 147, 427, 435
- NGC 185, 427, 435
- NGC 188, 443
- NGC 205, 426–27, 435
- NGC 253, 476, 479
- NGC 598, 471–72
- NGC 891, 479
- NGC 1052, 177
- NGC 1058, 227
- NGC 1068, 174, 183–84
- NGC 2685, 469
- NGC 2903, 476
- NGC 3031, 469, 473–74, 487
- NGC 3034, 469
- NGC 3079, 478
- NGC 3623, 468–69
- NGC 3718, 469
- NGC 3783, 181
- NGC 4051, 177
- NGC 4147, 434
- NGC 4151, 179–81, 183, 185
- NGC 4216, 468–69
- NGC 4258, 476
- NGC 4565, 468
- NGC 4590, 468–69
- NGC 4618, 228
- NGC 4631, 479
- NGC 4699, 223
- NGC 5055, 468, 476
- NGC 5128, 468–69
- NGC 5194, 472–73
- NGC 5236, 476, 488–89
- NGC 6717, 407
- NGC 6822, 426, 428
- NGC 6838, 452
- NGC 6946, 220, 474–75, 487
- NGC 7213, 177
- NGC 7331, 469
- NGC 7814, 468
- Nucleosynthesis
 - in massive stars, 329, 355–61, 363, 368
 - in standard model for Type I supernovae, 241–42, 246
 - in Type II supernovae, 211–12, 218, 223–26
- O
- O stars
 - and H II regions, 69, 79
 - interstellar reddening of, 15–16
 - mass loss from, 333, 337, 346, 352–53, 363–69
- OB associations, 54, 65, 68, 71, 80–81, 83, 369, 524–25
- OB stars, 54, 72, 329
 - mass loss from, 333–36, 346–51, 355
- Optical absorption-line studies
 - of interstellar gas, 499–535
 - instruments for, 508–11
- Orion Nebula, 16, 82
- Orion OB association, 518, 525
- Orion region, 525
- IRc2 in, 75
- Kleinmann-Low region in, 75, 82
- OSO 7 observations, 37
- OSO 8 observations, 35–38
- P
- P Cygni, 344
- Parametrized Post-Newtonian coordinate system (PPN), 543–53
- Photoelectric photometry, 1–3, 11–19
- Planetary nebulae, 50, 53, 64
- Polarization, 159–61, 167
 - in active galactic nuclei, 183–84
 - in galaxies, 460–61
 - optical observations of, 467–69
 - radio observations of, 469–78
 - in pulsars, 292–93, 307–11, 316, 322
- Population I disk stars
 - distribution of, 579–93
 - luminosity function of, 592–93, 601, 604
 - scale heights and scale lengths in, 595–96, 602
- Population I stars, 220–21. See also Stellar populations
 - main sequence, 443, 445
 - mass loss in, 380–81, 386, 390–95, 406
- Population II spheroidal stars, 579–602
 - axis ratio of, 597–98
 - distribution of, 579–94
 - luminosity function of, 593–94, 600
 - scale heights and scale lengths in, 595–96, 602
- Population II stars, 422, 426, 428–45. See also Stellar populations
 - main sequence, 437–46
 - mass loss from, 405–10
- Presupernova evolution, 206–12, 228–30, 347–48
- PSR 0525 + 21, 319
- PSR 0529 – 66, 288
- PSR 0531 + 21 (Crab Nebula pulsar), 293–95, 299, 318–19, 539
- PSR 0655 + 64, 301, 303–4, 565
- PSR 0809 + 74, 312–13
- PSR 0818 – 13, 312
- PSR 0820 + 02, 305–6
- PSR 0823 + 26, 299
- PSR 0826 – 34, 312

PSR 0833 - 45 (Vela X pulsar), 293-95, 299, 309, 318-19
 PSR 0950 + 08, 299, 310, 316
 PSR 1237 + 25, 570
 PSR 1509 - 58, 293-95, 299
 PSR 1855 + 09, 286, 304
 PSR 1913 + 16, 303-4, 537, 563-64, 567, 572-73
 PSR 1929 + 10, 299
 PSR 1937 + 21, 286, 288, 292, 303-7, 537, 540, 543, 545, 547-49, 553, 571
 PSR 1944 + 17, 312-13
 PSR 1953 + 29, 286, 304-7
 PSR 2303 + 46, 304
 Pulsar as coordinate time clock, 547-49
 Pulsar spin stability, 542-43
 Pulsar timing, 290-91, 537-75 and gravitational waves, 567-73
 Pulsars, 285-327, 479-80
 arrival-time of
 model of, 549-62
 observations of, 538-43
 in binary systems, 286, 292, 303-7, 323, 537, 554-67, 572-73
 birth and evolution of, 294-307
 birth rate of, 297, 302, 322
 distances of, 288, 299-300
 distribution of, 286-87, 322
 spatial, 300-1
 emission mechanism in, 307-21
 with interulses, 315-16, 322
 luminosity evolution of, 295-97, 302-3
 luminosity function of, 300-1
 magnetic fields in, 295, 297, 306, 321-22, 542
 magnetospheric theory of emission of, 316-19
 millisecond, 286, 288, 292, 298, 304-7, 316, 323, 543-45, 547-49, 553, 571
 noise in, 542-43, 547-48
 periods of, 288-91, 294-99, 322-23
 polarization in, 292-93, 307-11, 316, 322
 spectra of, 293-94
 subpulse drift and nulling in, 312-15
 and supernova remnants, 294-95, 303
 velocities of, 301-2
 waveforms of, 288-90

Q

Quasars, see also Quasi-stellar objects
 black holes in, 189-90
 broad-line region in, 185-91, 196, 198
 Quasi-stellar objects, 171-203
 emission-line region of, 177-81

R

R Andromedae, 96
 R Cygni, 94
 Radio astronomy, image restoration in, 127-70
 Radio-frequency emission, mass-loss determined from, 388-90
 Radio galaxies, 171-203
 broad-line region of, 185-98
 disk emission in, 198
 emission-line region of, 175, 177-85
 Radio sources, extragalactic, 479-80
 Rho Cassiopeiae, 413
 RR Lyrae variables, 424-26, 428-29, 433-37, 443, 446-52
 RS Canum Venaticorum systems, 400, 404-5, 414
 R 127, 344

S

S stars, 93-94, 96, 106-7, 114
 Sagittarius A, 172
 SC stars, 93, 106-7
 Sculptor system, 422, 426, 435
 Seyfert galaxies, 171-203. See also Active galactic nuclei, Active galaxies
 broad-line region of, 172-73, 177-98
 narrow-line region of, 173-76, 181-85, 198-200
 radial flow in, 174-75, 182-83, 191-94
 Seyfert 1 galaxies, 172-73, 177-84
 broad-line region in, 177-82
 narrow lines in, 181-82
 variations in spectra of, 179-80
 Seyfert 2 galaxies, 65, 68, 172-76, 182-85
 narrow-line region in, 173-76, 182
 Skylab observations, 25, 28, 35-37, 39, 395

Small Magellanic Cloud, 353, 365, 406, 426, 476-77
 SN 1006, 230
 SN 1885a, 247
 SN 1954a, 247
 SN 1957a, 233, 247
 SN 1961v, 227-28
 SN 19621, 245
 SN 19641, 245
 SN 19691, 220, 222
 SN 1972e, 236-37
 SN 1979c, 220
 SN 1980i, 247
 SN 1980k, 220
 SN 1981b, 236
 SN 1983k, 223
 SN 1983n, 245-46
 SN 1983v, 247
 SN 19841, 245
 SN 1985f, 228
 Solar atmosphere, 395-99
 abundances in, 398-99
 Solar atmosphere, transition region of, 23-48
 Doppler wavelength shifts in spectra of, 35-39
 dynamical energy balance models of, 42-46
 electron density in, 28-29
 empirical models of, 26-31
 jets in, 39-42
 loop model of, 43-46
 static energy balance models of, 31-32, 45
 temporal fluctuations in, 37-38
 turbulent events in, 39
 ultraviolet emission lines in, 23-28, 33-42
 Solar corona, 23-26, 33-35, 41, 44-46, 398-99, 414
 Solar coronal holes, 395-97, 399, 414
 Solar wind, 23, 26, 41, 395-99, 413-15
 Spicules, 26, 38-43, 46
 Spiral arms
 formation of, 459-60
 magnetic field alignment with, 468-69, 472-78
 Star-burst galaxies, 176
 Star counts, 349-50, 366-68, 577-611
 Star formation, 50, 72-76, 492
 Stars
 A-type, see A stars
 asymptotic giant branch, 405-7, 453
 B-type, see B stars
 binary, see Binary systems
 blue supergiant, 345-46
 carbon, see Carbon stars
 chromospheres of, 379-88

- cool, see Cool stars, G stars,
K stars, M stars
distribution of, in Galaxy,
578-83
dwarf, mass loss from, 395-
400
dwarf Population II, 428,
437-46
F-type, see F stars
G-type, see G stars
high-velocity, 422-25, 428,
432-38
hydrogen-ionizing flux in,
69-73, 80
intermediate white dwarf,
433-35
K-type, see K stars
luminous blue variable (Hubble-Sandage variables),
329, 344-45, 349, 357,
363-64
M-type, see M stars
magnitudes of, 16-18
main-sequence, 429-32, 437-
49, 453
mass loss in, 52, 65, 75,
117, 329-75, 377-420
massive, see Massive stars
metal-deficient, mass loss
from, 381, 405-10
metallicity of, 422-23, 432-
35
molecules in, 89-125
O-type, see O stars
OB, see OB stars
red giant, 93, 119, 347, 430.
See also K stars, M stars
red supergiant, 331-33, 345-
47, 353, 355, 364, 366-
67
mass-loss rates for, 337-
38
S-type, see S stars
subdwarf, 433-35, 441-45,
448, 454-55
supergiant, 329, 332-33,
344-46, 351-53, 366-
67
super-metal-rich, 20-22
syntrophic, 50, 405
velocity distribution of, 423-
25
Stellar atmospheres
abundances of molecules in,
92-96, 113
effect of molecules on struc-
ture of, 101-3, 118
inhomogeneity of, 113-15
Stellar evolution
accretion phase of, 52, 72, 75
and H II regions, 50-54, 65-
68
mass loss and, 329-75, 377-
420
HR diagram of globular clus-
ters and, 429-32
Stellar populations, 421-58. See
also Population I stars,
Population II stars
subsystems of, 437
Stellar winds
and H II regions, 68, 80, 82
and mass loss, 329, 333-38,
344-45, 348, 353, 358-
61, 368, 385, 387, 391-
94, 399, 408, 415
Sun
chromosphere of, 23-26, 37,
45
magnetic field of, 6, 24, 26,
28, 31, 44-46, 395
mass loss from, 378-79, 384,
386, 395-99
molecular spectra of, 111-12,
114-15
ultraviolet observations of,
395-97
ultraviolet spectra of, 23-42
nonthermal line broadening
in, 33-35
Sunspot spectra, molecules
observed in, 94, 96, 115
Supernova explosions, 54, 80,
205-53, 307, 358-61, 522
Supernova remnants
and H II regions, 52-53
and the interstellar medium,
503, 523-25
and pulsars, 294-95, 299,
303
Supernovae, evolution of, 205-
53
Supernovae, Type I, 206, 213,
228-47, 502, 524
evolution leading to, 228-30,
243-45
frequency of, 206
Hubble constant and, 243
nucleosynthesis in, 241-42,
246
peculiar, 245-47
standard model of, 234-42
thermonuclear models of,
230-42
Supernovae, Type II, 206-228,
524
"delayed" explosions of, 217-
19, 225
evolution of, 206-19, 226-27
core collapse in, 212-16
from massive stars, 226-27
frequency of, 206
light-curves of, 219-23
nucleosynthesis in, 223-26
rotation of, 219
subclasses of, 219-23
unusual, 227-28
T
T Tauri, 52
Time, 537-75
Time clock, pulsar as a coordi-
nate, 547-49
Time coordinate, relativistic,
543-49
Time scales, 539-42
TAI, 539-40, 544, 546-49
TDB, 539-40, 546-47
UTC, 539-40
TX Pisces, 94-95
U
Ultraviolet absorption-line stud-
ies of interstellar gas, 499-
535
instruments used for, 511-13
Ultraviolet observations
of active galactic nuclei, 180,
191
Ultraviolet spectroscopy, solar
Doppler shifts in, 35-37
nonthermal line broadening
in, 33-35
Ultraviolet spectrum, used for
mass-loss detection, 329,
378, 385-88, 390
UX Arietis, 400
V
Vela X, 293-95, 299, 309,
318-19, 524
VLA observations, 172, 184,
389-90, 400, 476, 478,
488
VLBI astrometry, 299
Voyager observations, 399
W
White dwarfs
in pulsar binary systems,
304-7
as supernovae precursors,
206-7, 229-36, 243-44,
246
Wolf-Rayet stars
abundances in, 357-58
emission nebulae around,
50
formation of, 363-65

galactic distribution of, 366-68
 mass loss in, 329, 342-47, 350, 353
 mass-loss rates in, 333-37, 342
 nucleosynthesis in, 355-61
 properties of, 361-63, 365-66
 spectra of, 361-62
 W49N, 76

XYZ

X-ray emission, 393-95, 524
 X-ray imaging, 132
 use of CCDs for, 282
 Y Canum Venaticorum, 95, 105
 YY Orionis, 52
 Zeeman effect, 6, 464
 Zero-age main sequence, 52-53, 65, 72-73, 349, 355, 369, 442-45, 447

Zeta-Aurigae/VV Cephei systems, 403-4

MISCELLANEOUS

22 Vulpeculae, 392
 47 Tucanae, 407, 430, 443, 449, 452, 594, 600
 I Zw 92, 175
 III Zw 77, 181